

# PROOF OF THE RIEMANN HYPOTHESIS

CHENG, YUAN-YOU FU-RUI

*Dedicated to my parents, my wife, and, my son*

## 1. INTRODUCTION

*If I were to awaken after having slept for five hundred years, my first question would be: Has the Riemann hypothesis been proven? [1]*

David Hilbert

Throughout this article, we shall use notations  $\mathbb{P}$  for the set of all prime numbers,  $\mathbb{P}^{\mathbb{N}}$  that of the all prime powers,  $\mathbb{N} = \{1, 2, 3, \dots\}$  that of natural numbers,  $\mathbb{Z}$  that of the integers,  $\mathbb{R}$  that of the real numbers, and  $\mathbb{C}$  that of the complex numbers. We adopt the notation  $a\mathbb{N} + b$ , to denote the subset of  $\mathbb{Z}$  containing all integers in the form of  $an + b$ , where  $n \in \mathbb{N}$  and  $a, b \in \mathbb{Z}$  are fixed. We also use the notation  $\mathbb{R}^+$  for the set of all positive real numbers. We shall use the symbol  $\epsilon \in \mathbb{R}^+$  for an arbitrary small positive real number, not necessarily the same at each occurrence in a given statement. We use the notation  $g(x) = f(x) + \text{Ie}(h(x))$  to represent the fact that  $|g(x) - f(x)| \leq h(x)$ . Suppose that  $g(x)$  and  $h(x)$  are complex functions of the variable  $x$  and  $f(y)$  is a positive real-valued function of  $y$ . The notation  $g(x) = h(x) + O(f(y))$  represents the fact that  $|g(x) - h(x)| \leq Bf(y)$  with some absolute constant  $B > 0$  whenever  $y$  is sufficiently large or  $y \geq y_0$  for some fixed positive number  $y_0$ . For convenience, we also use the notation  $f(x) = h(x) \oslash g(y)$  for the statement  $|f(x) - h(x)| \leq g(y)$ .

An arithmetic function  $f(n)$  is a complex-valued one defined for each  $n \in \mathbb{N}$ . The sum function  $F(x)$  for an arithmetic function  $f(n)$  is a piecewise constant function with possible discontinuities only at some  $n$ 's for  $n$  in a subset of  $\mathbb{N}$ . It is convenient for us to adopt the half-maximum convention for every such sum function; henceforth, we regard that every sum function

$$(1.1) \quad F(x) = \begin{cases} \frac{1}{2}f(x) + \sum_{n < x} f(n), & \text{if } x \in \mathbb{N}; \\ \sum_{n \leq x} f(n), & \text{otherwise.} \end{cases}$$

Therefore, in our work, every sum function for an arithmetic function  $f(n)$  satisfies the following property:

$$(1.2) \quad F(x) = \frac{F(x-0) + F(x+0)}{2},$$

---

2000 *Mathematics Subject Classification.* Primary 32A60, 11M26, 11N05.

at each point  $x$  where  $F(x-0)$  and  $F(x+0)$  are the left sided limit and right sided limit of  $F(x)$  at  $x$ , respectively. We also use the Heaviside function  $H_\alpha(v)$  at any fixed point  $\alpha \in \mathbb{R}$ , which is defined by

$$(1.3) \quad H_\alpha(v) = \begin{cases} 0, & \text{for } 0 < v < \alpha, \\ \frac{1}{2}, & \text{for } v = \alpha, \\ 1, & \text{for } v > \alpha. \end{cases}$$

It is obvious that prime numbers play a central role in number theory. It has been understood that in-depth study of the distribution of primes is connected to the Riemann zeta function since 1859, by Riemann's epoch-making work [30]. The Riemann zeta function  $\zeta(s)$  is a regular complex-valued function for  $s \in \mathbb{C} \setminus \{1\}$ . We shall accustomedly denote  $s = \sigma + it$ . For  $\sigma > 1$ , the Riemann zeta function is defined by

$$(1.4) \quad \zeta(s) = \sum_{n=1}^{\infty} \frac{1}{n^s} = \prod_{p \in \mathbb{P}} \frac{1}{1 - \frac{1}{p^s}},$$

in which the last equation was known for a real variable  $s$  by Leonard Euler as early as 1773, and may be regarded as an analytic version of the fundamental theorem of arithmetic, which states that for each natural number  $n$  there is a unique factorization

$$(1.5) \quad n = \prod_{p^a \parallel n} p^a,$$

where  $p^a \parallel n$  means that  $a$  is the highest positive exponent such that  $p^a$  divides  $n$ .

The definition in (1.4) may be extended to the whole complex plane by many different ways. For references, one may see [1], [9], [16], or [22]. For  $\sigma > 0$ , we have

$$(1.6) \quad \zeta(s) = \frac{s}{s-1} - s \int_1^\infty \frac{v - [v]}{v^{s+1}} dv \quad \text{for } \sigma > 0,$$

where  $[v]$  is the integer part of  $v$  or the greatest integer less than or equal to  $v$ . One may notice that  $s = 1$  is a pole for  $\zeta(s)$  having residue 1. For  $s \in \mathbb{C} \setminus \{1\}$ , one may use the functional equation

$$(1.7) \quad \pi^{-\frac{s}{2}} \Gamma\left(\frac{s}{2}\right) \zeta(s) = \pi^{-\frac{1-s}{2}} \Gamma\left(\frac{1-s}{2}\right) \zeta(1-s),$$

to consider the zeros of the Riemann zeta function. Here,  $\Gamma(s)$  is a complex-valued function extending the definition for the factorial  $n!$  such that  $\Gamma(n+1) = n!$ . One may define it by

$$(1.8) \quad \Gamma(s) = \frac{1}{s} e^{\gamma_0 s} \prod_{n=1}^{\infty} \frac{e^{s/n}}{1 + s/n},$$

where  $\gamma_0 \approx .577215665 \dots$  is Euler's constant. The Gamma function  $\Gamma(s)$  has simple poles for  $s \in -\mathbb{N} + 1$  and no zeros at all; in order to compensate for all poles of  $\Gamma(s)$ , except for that at  $s = 0$ ,  $\zeta(s)$  vanishes at  $s \in -2\mathbb{N}$ , while the pole of  $\zeta(s)$  at  $s = 1$  corresponds the pole of  $\Gamma(s)$  at  $s = 0$ . These

zeros  $s \in -2\mathbb{N}$  for  $\zeta(s)$  are called trivial zeros; all other zeros are called non-trivial zeros. The identity

$$(1.9) \quad \zeta(\bar{s}) = \overline{\zeta(s)},$$

from the reflection principle in complex analysis shows that all non-trivial zeros are symmetric about the real line. From the functional equation (1.7), one sees that all non-trivial zeros are symmetric about the line  $\sigma = \frac{1}{2}$  even though  $\zeta(s)$  is not symmetric about the line  $\sigma = \frac{1}{2}$ . It is proved that all non-trivial zeros for the Riemann zeta function are located in the so-called critical strip  $0 \leq \sigma \leq 1$ .

The **Riemann hypothesis** states there is no zeros for  $\zeta(s)$  for  $\sigma > \frac{1}{2}$ , which is equivalent to the statement that all the non-trivial zeros are located on the line  $\sigma = \frac{1}{2}$  by the property of symmetry from its functional equation. This conjecture was made by Riemann in 1860 together with six other conjectures in [30]. Every other conjecture made by Riemann has been proved since that time, but the Riemann hypothesis has resisted every attack so far.

It was proved in 1890's by Hadamard and de la Vallée Poussin that  $\zeta(s)$  does not have zeros on the line  $\sigma = 1$ . However, we are unaware as to whether the Riemann zeta function does not have zeros for  $\sigma > a$  with any fixed  $a < 1$  at the time of this work being written. The best known zero free region so far, from Vinogradov's method, was in the form of

$$(1.10) \quad \sigma > 1 - \frac{1}{49.13 (\log |t|)^{2/3} (\log \log |t|)^{1/3}},$$

see [18].

In another direction, we denote the number of zeros of  $\zeta(s)$  on the critical line  $\frac{1}{2} + it$  such that  $0 < t < T$  by  $M(T)$ . Related to this, Hardy proved in 1941 that there are infinitely many zeros of  $\zeta(s)$  on the critical line. Selberg improved this in 1942. He proved that  $M(T) > C \frac{T}{2\pi} \log \frac{T}{2\pi}$  for some constant  $C$  for  $T > T_0$  for some  $T_0 > 0$  with  $0 < C < 1$ . Also, Levinson showed in 1974 that at least a third of nontrivial zeros of  $\zeta(s)$  are on the critical line. This has been improved to two-fifths by Conrey in 1989. For references, one may see [17], [33], [26], and [14].

On the computation side, we mention here that Gourdon calculates the first  $10^{13}$  zeros of  $\zeta(s)$  on the critical line in 2004, see [20]. Such computations involve Gram's law, Rosser's rule, Turing's method, and the Odlyzko-Schönhage Algorithm which is a heuristics for the location of zeros on the critical line, see [3], [36], and [28].

## 2. THE MAIN RESULTS

*If you could be the Devil and offer a mathematician to sell his soul for the proof of one theorem - what theorem would most mathematicians ask for? I think it would be the Riemann hypothesis. [25]*

Hugh L. Montgomery

Perhaps, one should say that it is important to consider the Riemann hypothesis in terms of how it relates to number theory, as its principal application concerns the prime numbers. The prime indicator function,  $\text{Id}_{\mathbb{P}}(n)$  for every  $n \in \mathbb{N}$ , is defined by

$$(2.1) \quad \text{Id}_{\mathbb{P}}(n) = \begin{cases} 1, & \text{if } n \text{ is a prime number;} \\ 0, & \text{if } n \text{ is not a prime number.} \end{cases}$$

The prime counting function  $\pi(x)$ , or the sum function for the prime indicator, is defined by

$$(2.2) \quad \pi(x) = \sum_{n \leq x} \text{Id}_{\mathbb{P}}(n) = \begin{cases} \frac{1}{2} + \sum_{n < x} \text{Id}_{\mathbb{P}}(n), & \text{if } x \in \mathbb{P}; \\ \sum_{n \leq x} \text{Id}_{\mathbb{P}}(n), & \text{otherwise,} \end{cases}$$

with the notation  $\sum$  meaning that the half-maximum convention is adopted. Gauss' logarithmic integral function  $\text{Li}(x)$  is defined by

$$(2.3) \quad \text{Li}(x) = \int_2^x \frac{du}{\log u}.$$

The *prime number theorem* in its best known form is

$$(2.4) \quad \pi(x) = \text{Li}(x) + O\left(x \exp\left(-C(\log x)^{3/5}(\log \log x)^{-1/5}\right)\right),$$

where  $C$  is positive constant. Let  $0 < \lambda < 1$  and  $T > 0$ . The *density hypothesis* says that

$$(2.5) \quad N(\lambda, T) = O(T^{2(1-\lambda)+\epsilon}),$$

where  $N(\lambda, T)$  is the number of zeros for  $\zeta(s)$  in the region for which  $\sigma \geq \lambda$  and  $0 < t \leq T$ . It is easy to see that  $N(\lambda, T) = 0$  when  $\lambda > 1$  from Euler's product formula, the second equation in (1.4). It was proved independently by Hadamard and de la Vallée Poussin in 1896 that  $N(1, T) = 0$ . In the case that  $\lambda = 0$ , it has been known, since 1905, by means of the Riemann-von Mangoldt Theorem that

$$(2.6) \quad N(T) := N(0, T) = \frac{T}{2\pi} \log \frac{T}{2\pi} + O(\log T).$$

We shall call the following statement *the strong density hypothesis*:

$$(2.7) \quad N(\lambda, T) = O(T^{2(1-\lambda)} \log T).$$

It is well-known that the Riemann hypothesis implies the Lindelöf hypothesis and the Lindelöf hypothesis implies the density hypothesis. However, it has not been known whether the density hypothesis implies the Riemann hypothesis. In this article, we prove the following result.

**Theorem 1.** *The strong density hypothesis implies the Riemann hypothesis.*

In the writing process of this paper, we actually proved an even stronger result as follows, see [11]. We have,

$$(2.8) \quad N(\lambda, T) = O(\log T),$$

for all  $\frac{1}{2} < \lambda < 1$ . Therefore, we have the following theorems.

**Theorem 2.** *The Riemann hypothesis is valid; i.e., all non-trivial zeros of the Riemann zeta function lie on the line  $s = \frac{1}{2}$ .*

The Riemann hypothesis is the analytic version for the prime number theorem in its certain form in number theory. The Riemann hypothesis is equivalent to the assertion on the prime counting in the form of (2.9) as follows, which is stated in [15] and is from [34] with some calculation for the smaller numbers.

**Theorem 3.** *The prime number theorem in the following form holds. That is,*

$$(2.9) \quad \pi(x) = \text{Li}(x) \trianglelefteq \frac{1}{8\pi} x^{\frac{1}{2}} \log x,$$

for  $x \geq 1451$ . The coefficient  $\frac{1}{8\pi}$  may be replaced by 1 if  $x \geq 2.01$ .

We actually can obtain stronger result with the information on the differences of non-trivial zeros of the Riemann zeta function, see [10].

**Theorem 4.** *The Lindelöf hypothesis is valid; that is,*

$$(2.10) \quad \zeta\left(\frac{1}{2} + it\right) = O(t^\epsilon).$$

In [8], it is proved by applying van der Corput's method in 1921 that in (2.10)  $\epsilon = \frac{1}{6} + \frac{\log \log t}{\log t}$  with the constant  $B = 3$  instead of the  $O$  notation. The best known result in this direction is slightly better than  $\frac{1}{6}$ . The validity of Theorem 4 from the Riemann hypothesis is well-known. We leave to reader on whether Linderlöf hypothesis implies the strong density hypothesis defined in (2.7) or the even stronger result in (2.8).

We define the function  $\psi(x)$ , as in the literature, and the  $\varpi$ -function  $\varpi(x)$  by

$$(2.11) \quad \psi(x) = \sum_{n \leq x} \Lambda(n), \quad \varpi(x) = \sum_{n \leq x} (\Lambda(n) - 1),$$

where

$$(2.12) \quad \Lambda(n) = \begin{cases} \log p, & \text{if } n = p^k \in \mathbb{P}^{\mathbb{N}}, \\ 0, & \text{otherwise.} \end{cases}$$

We shall prove the following Theorem 5. From (2.11), it is straightforward to see that (2.13) is equivalent to the statement  $\psi(x) = x + O(x^{1/2} \log^2 x)$ . Also, it is a standard result in the literature that  $\psi(x) = x + O(x^{1/2} \log^2 x)$  is equivalent to Theorem 2 and Theorem 3. Therefore, Theorem 5 is equivalent to Theorem 1.

**Theorem 5.** *The strong density hypothesis implies*

$$(2.13) \quad \varpi(x) = O(x^{1/2} \log^2 x).$$

## 3. PROOF OF THEOREM 5

*It would be very discouraging if somewhere down the line you could ask a computer if the Riemann hypothesis is correct and it said, “Yes, it is true, but you won’t be able to understand the proof.”*

Ronald L. Graham

Corresponding to (2.11), we study the function

$$(3.1) \quad \mathcal{Z}(s) = -\frac{\zeta'(s)}{\zeta(s)} - \zeta(s) = \sum_{n=1}^{\infty} \frac{\Lambda(n) - 1}{n^s}.$$

The definition in (3.1) is valid for  $\sigma > 1$  because the series in (3.1) converges absolutely and uniformly in the half-plane  $\sigma \geq 1 + \epsilon$ . Actually, we shall validify the definition in (3.1) for  $\sigma > \frac{1}{2}$  during the process of proving Theorem 5.

For convenience, we define

$$(3.2) \quad H_0(x) = \begin{cases} 1 - \frac{1}{57 \log^{2/5} x (\log \log x)^{1/5}} - \frac{7 \log \log x}{5 \log x}, & x > x_0, \\ \frac{1}{2}, & x \leq x_0, \end{cases}$$

with  $x_0 \approx 46.784$ . We then define

$$(3.3) \quad \begin{cases} H_j = H_{j-1} - \frac{\log \log x}{8 \log x}, & x > x_j, \\ \frac{1}{2}, & x \leq x_j, \end{cases}$$

where  $x_j$  is the solution of  $H_j(x) = \frac{1}{2}$ , for all  $j \in \mathbb{N}$ . It is obvious that  $x_j > x_{j-1}$  and  $H_j(x) > H_{j-1}(x)$  for  $x > x_j$ .

Recall the Euler product formula as the second equation in (1.4). It is well-known that the Riemann zeta function does not vanish for  $\sigma > 1$  from the Euler product formula. Also, recall the zero-free region described in (1.10), from which, one can show that

$$(3.4) \quad \varpi(x) = O(x^{H(x)} \log^2 x),$$

when  $H(x) = H_0(x)$  if only we allow the constant in the  $O$ -notation to be large enough. For references, one may see [13]. Because of the same understanding, we may assume that this is valid for  $x \leq 96726620$ . From now on, let  $y = \frac{3}{4}x$  and  $T = \log y$  so that  $y \geq 72544965$  and  $T > 18$ . Besides, we assume that the constant for the  $O$ -notation in (3.4) is fixed henceforth.

Under the assumption of Theorem 5, we claim here that (3.4) is valid when  $H(x) = H_j(x)$  only if (3.4) is true when  $H(x) = H_{j-1}$ , for every  $j \in \mathbb{N}$ . From the start that (3.4) is valid when  $H(x) = H_0(x)$ , it is straightforward that (3.4) is justified when  $H(x) = H_j(x)$  for all  $j \in \mathbb{N}$  by induction. Obviously, the last statement means that (3.4) is valid for  $H(x) = \frac{1}{2}$ . Therefore, our claim is equivalent to Theorem 5.

What remains is to prove our claim in this section, with details being put in sections afterwards.

First, we recall the definition the Heaviside function  $H_1(v)$  in (1.3). Let  $m = h(T) + \frac{\log \log y}{8 \log y}$  and  $\delta_T(v)$  and  $\delta_E(v)$  be defined in (5.2) and (5.3) with respect to the parameter  $m$ , respectively. Then

$$(3.5) \quad H_1(v) = \delta_T(v) + \delta_E(v).$$

Next, we recall the definition of the  $\varpi$ -function  $\varpi(x)$  in (2.11). It follows that

$$(3.6) \quad \varpi(x) = \sum_{n \leq x} (\Lambda(n) - 1) = \sum_{n=1}^{\infty} H_1\left(\frac{x}{n}\right) (\Lambda(n) - 1) = \varpi_1(x) + \varpi_2(x),$$

where  $\varpi_1$  and  $\varpi_2$  are defined by

$$(3.7) \quad \varpi_1(x) = \sum_{n=1}^{\infty} \delta_T\left(\frac{x}{n}\right) (\Lambda(n) - 1), \quad \varpi_2(x) = \sum_{n=1}^{\infty} \delta_E\left(\frac{x}{n}\right) (\Lambda(n) - 1).$$

Now, we consider the differences  $\varpi_1(x) - \varpi_1(y)$  and  $\varpi_2(x) - \varpi_2(y)$  in preparing for the induction with respect to  $x$  under the assumption  $0 < y < x$ . In Section 5, we acquire

$$(3.8) \quad \begin{aligned} \varpi_1(x) - \varpi_1(y) &= O\left(\frac{(x-y)y^m T^{3/2}}{y}\right) \\ &\quad + \frac{2A(x-y)T^{2[1-h(T)]} \log y \log T}{y^{1-h(T)} (\log y - 2 \log T)}. \end{aligned}$$

In Section 6, we obtain

$$\varpi_2(x) - \varpi_2(y) = \frac{1}{2\pi i} \int_{\mathcal{L}} \mathcal{Z}(s) \frac{x^s - y^s}{s} ds + O(\log x).$$

??, therefore,

$$(3.9) \quad \varpi_2(x) - \varpi_2(y) = O\left(\frac{2A(x-y)y^m T \log T}{y}\right).$$

Note that  $h(T) \geq \frac{1}{2}$  and  $\frac{\log y}{\log y - 2 \log T} \cdot \frac{\log \log y}{\log^{1/2} y} \leq 1.001$ . It follows that

$$(3.10) \quad \begin{aligned} \varpi(x) - \varpi(y) &= O\left(\frac{(x-y)y^m \log^{3/2} y}{y}\right) \\ &\quad + O\left(\frac{(x-y)y^{h(T)} \log \log y}{y}\right) + O\left(\frac{(x-y)y^m \log y \log \log y}{y}\right) \\ &= O\left(\frac{(x-y)y^{\min\{m, 1/2\}} \log^{3/2} y}{y}\right). \end{aligned}$$

Recalling the definition of  $m$ , we see that the constant in the last  $O$  notation being only very slightly bigger than that in the first  $O$  notation.

We may inductively assume that (3.4)? is valid with  $x$  being replaced by  $y$ . Note that  $H(x) - H(y) \geq x^{1/2} - y^{1/2} = \frac{16}{7}x^{1/2} \geq 1$ . Note that  $\log y \geq \frac{2(x-y)}{y \log^{1/4} y}$  from the designation of  $y$  and  $\frac{2(x-y)}{y \log^{1/4} y} > \log\left(1 + \frac{x-y}{y \log^{1/4} y}\right)$

from  $2w > \log(1+w)$  for any  $0 < w < 1$ . It follows that  $H(x) \log x + \frac{7}{4} \log \log x - H(y) \log y - \frac{7}{4} \log \log y > \log\left(1 + \frac{x-y}{y \log^{1/4} y}\right)$ . Therefore, we have

$$(3.11) \quad \frac{x^{H(x)} \log^{7/4} x}{y^{H(y)} \log^{7/4} y} > 1 + \frac{x-y}{y \log^{1/4} y}.$$

Combining (3.10) and (3.4)? with  $x$  being replaced by  $y$  and utilizing (3.11), we conclude that (3.4)? is valid.

To finish the proof of the proposition, we let  $x = x_l$ ,  $x_{l-1} = \frac{3}{4}x_l$ ,  $\dots$ ,  $x_1 = \frac{3}{4}x_2$ ,  $x_0 = \frac{3}{4}x_1$  for sufficiently large  $l$  so that  $x_0 < 96726620 \log x_1$ . We conclude that the proposition is valid by induction.

#### 4. THE FUNCTION DEFINED BY THE SERIES

*There have probably been very few attempts at proving the Riemann hypothesis, because, simply, no one has ever had any really good idea for how to go about it! [4]*

Atle Selberg

It is not clear whether the series on the right side of (3.1) is convergent for  $\frac{1}{2} < \sigma \leq 1$ . The affirmative answer implies the Riemann hypothesis.

It is well-known that the remainder term of the prime number theorem may be derived from the zero-free region of the Riemann zeta function. Less known is that converse is also true, see Turán's work [35] in 1950. Following Turán's power sum method, we prove the following lemma in [6]. (The exact definition of  $h_j(t)$  will be made after we finish that paper.)

**Lemma 1.** *Let  $H_j(x)$  for  $j \in \{0\} \cap \mathbb{N}$  be defined as in (3.2) and (3.3). Suppose that  $H_j(x)$  is suitable for (3.4) to be valid for a  $j \in \{0\} \cap \mathbb{N}$ . Then, there is a corresponding function  $h_j(t)$  such that  $\frac{1}{2} \leq h_j(t) < 1$ ,  $\lim_{j \rightarrow \infty} h_j(t) = \frac{1}{2}$ , and  $\mathcal{Z}(s)$  is defined by*

$$(4.1) \quad \mathcal{Z}(s) = \sum_{n=1}^{\infty} \frac{\Lambda(n) - 1}{n^s},$$

for  $\sigma > h_j(t)$ . Also, both the functions  $\mathcal{Z}(s)$  and  $\zeta(s)$  are regular and  $\zeta(s)$  does not vanish in the same region.

Here, we instead give a simple idea on why Lemma 1 appears to be valid. By the partial summation method, we acquire

$$(4.2) \quad \begin{aligned} \mathcal{Z}(s) &= \sum_{n=1}^{\infty} \frac{\Lambda(n) - 1}{n^s} = \sum_{n=1}^{\infty} \varpi(n) \left( \frac{1}{n^s} - \frac{1}{(n+1)^s} \right) \\ &= s \sum_{n=1}^{\infty} \varpi(n) \int_n^{n+1} \frac{du}{u^{s+1}} = s \int_1^{\infty} \frac{\varpi([u])}{u^{s+1}} du. \end{aligned}$$



Note that  $\varpi(\lfloor u \rfloor) = \varpi(u) - (\varpi(u) - \varpi(\lfloor u \rfloor)) = \varpi(u) - \frac{1}{2}(\Lambda(\lfloor u \rfloor) - 1)$ . One sees

$$(4.3) \quad \mathcal{Z}(s) = s \int_1^\infty \frac{\varpi(u)}{u^{s+1}} \, du - \frac{s}{2} \int_1^\infty \frac{\Lambda(\lfloor u \rfloor) - 1}{u^{s+1}} \, du.$$

Denote

$$(4.4) \quad \varrho(u) = \int_1^u \varpi(v) \, dv.$$

Recalling the definition of  $\varrho(u)$  in (4.4) and applying the integration by parts, we get

$$(4.5) \quad \begin{aligned} \int_1^\infty \frac{\varpi(u)}{u^{s+1}} \, du &= \int_1^\infty \frac{d\varrho(u)}{u^{s+1}} \\ &= \lim_{u \rightarrow \infty} \frac{\varrho(u)}{u^{s+1}} + (s+1) \int_1^\infty \frac{\varrho(u)}{u^{s+2}} \, du, \end{aligned}$$

since  $\varrho(1) = 0$ . Combining (4.3) and (4.5) and recalling the definition of  $\varrho(u)$  in (4.4), we obtain

$$(4.6) \quad \mathcal{Z}(s) = s \lim_{u \rightarrow \infty} \frac{\varrho(u)}{u^{s+1}} + s(s+1) \int_1^\infty \frac{\varrho(u)}{u^{s+2}} \, du - \frac{s}{2} \int_1^\infty \frac{\Lambda(\lfloor u \rfloor) - 1}{u^{s+1}} \, du.$$

The last integral converges absolutely and uniformly for  $\sigma \geq \epsilon$  as  $\Lambda(\lfloor u \rfloor) - 1 \ll \log u$ . Note that  $\varrho(u) = O(u^{H(u)+1} \log^2 u)$  from the assumption  $\varpi(u) = O(u^{H(u)} \log^2 u)$ . Suppose that  $\sigma > H(u)$ . Then, the limit in (4.6) is 0. Also, the first integral in (4.6) converges absolutely and uniformly for  $\sigma \geq H(u) + \epsilon$ . Therefore, the function is defined by the first series in (4.2). By Weierstrass M-test, we see that  $\mathcal{Z}(s)$  is a regular function for  $\sigma > H(n)$ .

From this argument, we see that the function  $\mathcal{Z}(s)$  is defined by the series in (4.6) so that it is a regular function which does not vanish for  $\sigma > h(t)$ , for some function  $h(t)$ . It follows that the function  $\mathcal{Z}(s)$  does not have poles and is regular for  $\sigma > h(t)$ . Therefore, the function  $-\frac{\zeta'(s)}{\zeta(s)}$  does not have poles in the region. However, the last statement means that  $\zeta(s)$  does not vanish in the region.

We end this section by stating the upper bound of  $\zeta(s)$ ,  $-\frac{\zeta'(s)}{\zeta(s)}$ , and  $\mathcal{Z}(s)$ . From [22], we recall

$$(4.7) \quad |\zeta(s)| = O(|t|^{3/2}), \quad \sigma \geq -1.$$

We also recall Proposition 9.4 in [12] with  $|\frac{\zeta'(s)}{\zeta(s)}| = O(\log^2 |t|)$  for  $t \geq 12$  and  $-1 \leq \sigma \leq 2$ . Noting that  $|\frac{\zeta'(s)}{\zeta(s)}| = O(1)$  for  $0 \leq t < 12$  and  $-1 \leq \sigma \leq 2$ , we see that

$$(4.8) \quad |\mathcal{Z}(s)| = O(|t|^{3/2}), \quad -1 \leq \sigma \leq 2.$$

## 5. APPLYING AN INTEGRAL TRANSFORM FORMULA

*A science is said to be useful if its development tends to accentuate the existing inequalities in the distribution of wealth or more directly promotes the destruction of human life. The theory of prime numbers satisfies no such criteria. [1]*

Godfrey H. Hardy

It is known that there are no zeros for the Riemann zeta function for  $\Im(s) \leq 14$  in the critical strip. Henceforth, we assume that  $T \geq 14$  and  $T \leq T_a < T + 1$  such that  $T_a \neq \gamma$  for any zero  $\rho = \beta + i\gamma \in \mathbf{Z}$ .

The classical formula

$$(5.1) \quad \frac{1}{2\pi i} \int_{m-i\infty}^{m+i\infty} \frac{v^s}{s} \, ds = H_1(v),$$

where  $H_1(v)$  is the step function defined in (1.3), plays an important role in the studies of the prime numbers. However, it is used only for  $m > 1$  in the literature. It is crucial for us to use this formula for  $\frac{1}{2} < m \leq 1$ . By designation in Section 3, we have  $N(m, T + 1) = 0$  because . This is justified in Section 4. We do not need the formula in (5.1), instead, we shall use the decomposition of the Heaviside function  $H_1(v)$  in (3.5), with  $\delta_T(v)$  and  $\delta_E(v)$  being defined as follows. Let  $\mathcal{S}$  be the route along the vertical line segment from  $m - iT_a$  to  $m + iT_a$ . We denote

$$(5.2) \quad \delta_T(v) := \frac{1}{2\pi i} \int_{\mathcal{S}} \frac{v^s}{s} \, ds, \quad m > \frac{1}{2}.$$

Then, let  $\mathcal{L}$  be a simple curve from  $m + iT_a$  to  $m - iT_a$  such that the union  $\mathcal{L}'$  of  $\mathcal{S}$  and  $\mathcal{L}$  is a simply closed curve and the index of the point  $s = 0$  with respect to  $\mathcal{L}'$  is equal to 1. Also, let  $\mathcal{R}$  be a simple curve from  $m + iT_a$  to  $m - iT_a$  such that the union  $\mathcal{R}'$  of  $\mathcal{S}$  and  $\mathcal{R}$  is a simply closed curve and the index of the point  $s = 0$  with respect to  $\mathcal{R}'$  is equal to 0. We also assume  $\mathcal{L} \cap \mathcal{R} = \emptyset$  so that  $\mathcal{L}' \cap \mathcal{R}' = \mathcal{S}$ . Note that the union of  $-\mathcal{R}$  and  $\mathcal{L}$  is a simply closed curve, where  $-\mathcal{R}$  is the route along  $\mathcal{R}$  with the opposite direction. We define

$$(5.3) \quad \delta_E(v) := \begin{cases} \frac{1}{2\pi i} \int_{\mathcal{R}} \frac{v^s}{s} \, ds, & 0 < v < 1, \\ \frac{1}{\pi} \int_{T_a/m}^{\infty} \frac{du}{1+u^2}, & v = 1, \\ \frac{1}{2\pi i} \int_{\mathcal{L}} \frac{v^s}{s} \, ds, & v > 1. \end{cases}$$

Note that  $s = 0$  is the unique pole of the function  $\frac{v^s}{s}$ , at which the function has a residue 1, in the whole complex plane. In the case  $v \neq 1$ , we justify (3.5) by the Argument Principle, noting that the route direction of  $\mathcal{R}'$  is clockwise. The equality (3.5) in the case  $v = 1$  is also valid. These may be done similarly to that in [16] or [22].

We then apply the Cauchy's Residue Theorem on the function  $\mathcal{Z}(s) \frac{x^s}{s}$ . There is no pole at  $s = 1$  for  $\mathcal{Z}(s)$  since the pole of  $-\frac{\zeta'(s)}{\zeta(s)}$  and that of  $\zeta(s)$

at  $s = 1$  is canceled. Recall

$$(5.4) \quad \begin{aligned} \frac{\zeta'(s)}{\zeta(s)} &= -\frac{1}{s-1} + \sum_{\rho \in \mathbb{Z}} \left( \frac{1}{s-\rho} + \frac{1}{\rho} \right) \\ &\quad + \sum_{n=1}^{\infty} \left( \frac{1}{s+2n} - \frac{1}{2n} \right) + \log(2\pi) - 1, \end{aligned}$$

which may be found in any standard references. We see that the residue of  $\frac{x^s}{s} \mathcal{Z}(s)$ , or that of  $-\frac{x^s}{s} \frac{\zeta'(s)}{\zeta(s)}$ , at each point  $s = \rho$  is  $-\frac{x^\rho}{\rho}$ . Using (5.4), we see that  $-\frac{\zeta'(0)}{\zeta(0)} = -\log(2\pi)$ . Also, recall (1.6) and the functional equation for the Riemann zeta function in the following form:

$$(5.5) \quad \zeta(s) = 2^s \pi^{s-1} \sin \frac{\pi s}{2} \Gamma(1-s) \zeta(1-s).$$

From these two equations, one may show that  $\zeta(0) = -\frac{1}{2}$  as in [29]. It follows that the residue of  $\frac{x^s}{s} \mathcal{Z}(s)$  at  $s = 0$  is equal to  $-\frac{\zeta'(0)}{\zeta(0)} - \zeta(0) = -\log(2\pi) + \frac{1}{2}$ . Now, recall the definition of  $\varpi_1(x)$  in (3.7). We have

$$(5.6) \quad \varpi_1(x) = \int_{m-iT_a}^{m+iT_a} \frac{x^s}{s} \sum_{n=1}^{\infty} \frac{\Lambda(n) - 1}{n^s},$$

by Cauchy's Residue Theorem.

Note here that (4.1) is valid. Let  $L_b$  and  $L_u$  be the horizontal line routes from  $m - iT_a$  to  $-1 - iT_a$  and from  $-1 + iT_a$  to  $m + iT_a$ , respectively, and,  $L_l$  be the vertical line route from  $-1 - iT_a$  to  $-1 + iT_a$ . By Cauchy's Residue Theorem, one gets

$$(5.7) \quad \begin{aligned} \varpi_1(x) &= \int_{m-iT_a}^{m+iT_a} \frac{x^s}{s} \mathcal{Z}(s) \, ds = \int_{m-iT_a}^{m+iT_a} \frac{x^s}{s} \left( -\frac{\zeta'(s)}{\zeta(s)} - \zeta(s) \right) \, ds \\ &= - \sum_{\rho \in \mathbb{Z}: |\Im(\rho)| \leq T_a} \frac{x^\rho}{\rho} + \frac{1}{2\pi i} \int_{L_b+L_l+L_u} \frac{x^s}{s} \mathcal{Z}(s) \, ds - \log(2\pi) + \frac{1}{2}, \end{aligned}$$

It follows that

$$(5.8) \quad \begin{aligned} \varpi_1(x) - \varpi_1(y) &= - \sum_{|\Im(\rho)| \leq T_a} \frac{x^\rho - y^\rho}{\rho} \\ &\quad + \frac{1}{2\pi i} \int_{L_b+L_l+L_u} \frac{\mathcal{Z}(s)(x^s - y^s)}{s} \, ds. \end{aligned}$$

We estimate the integrals in (5.8). First, the difference for the integral along with  $L_l$  is comparatively smaller and may be accessed directly without involving the difference with respect to  $x$  and  $y$ . For both  $w = x$  and  $y$ , we have

$$\int_{L_l} \frac{\mathcal{Z}(s)w^s}{s} \, ds = O(T^{3/2}) y^{-1} \int_{L_l} \, dt = O\left(\frac{T^{5/2}}{y}\right),$$

recalling (4.8) and  $y < x$ . We use  $s = -1 + it$  and  $|x^s| \leq x^{-1} < y^{-1}$  and  $|y_s| \leq y^{-1}$ . It follows that

$$(5.9) \quad \left| \frac{1}{2\pi i} \int_{L_l} \frac{\mathcal{Z}(s)(x^s - y^s)}{s} \mathrm{d}s \right| = O\left(\frac{T^{5/2}}{y}\right).$$

Note that if  $\sigma < 1$  then

$$(5.10) \quad \left| \frac{x^s - y^s}{s} \right| \leq \int_y^x |v^{s-1}| \mathrm{d}v = \int_y^x v^{\sigma-1} \mathrm{d}v \leq (x - y) y^{\sigma-1}.$$

Using this inequality and (4.8), we acquire

$$(5.11) \quad \frac{1}{2\pi i} \int_{L_b+L_u} \frac{\mathcal{Z}(s)(x^s - y^s)}{s} \mathrm{d}s = O\left(\frac{(x - y) T^{3/2}}{y^{1-m}}\right).$$

To estimate the sum in (5.8), we first note that

$$y^\beta = y^{1/2} + \log y \int_{1/2}^\beta y^v \mathrm{d}v,$$

and

$$\int_{1/2}^\beta y^v \mathrm{d}v = \int_{1/2}^1 y^v (1 - H_\beta(v)) \mathrm{d}v,$$

for each non-trivial zero  $\rho = \beta + i\gamma$  of the Riemann zeta function and  $H_\beta(u)$  is the Heaviside function at the point  $\beta$  with the half-maximum convention in (1.3). Recall that the Riemann zeta function does not have zeros for  $\sigma > h(t)$ . We also use (5.10) with  $s = \rho$  for each  $\rho \in \mathbb{Z}$ . It follows that

$$\begin{aligned} \left| \sum_{\rho \in \mathbb{Z}: |\gamma| \leq T_a} \frac{x^\rho - y^\rho}{\rho} \right| &\leq \frac{x - y}{y} \sum_{\rho \in \mathbb{Z}: |\gamma| \leq T_a} x^\beta \\ &\leq \frac{x - y}{y} \sum_{\rho \in \mathbb{Z}: |\gamma| \leq T_a} \left( y^{1/2} + \log y \int_{1/2}^1 y^v (1 - H_\beta(v)) \mathrm{d}v \right) \\ &= \frac{x - y}{y} \left[ y^{1/2} \sum_{\rho \in \mathbb{Z}: |\gamma| \leq T_a} 1 + \log y \int_{1/2}^{h(T)} y^v \sum_{\rho \in \mathbb{Z}: |\gamma| \leq T_a} (1 - H_\beta(v)) \mathrm{d}v \right]. \end{aligned}$$

Note that  $\sum_{\rho \in \mathbb{Z}: |\gamma| \leq T_a} 1 = N(T_a)$  and  $\sum_{\rho \in \mathbb{Z}: |\gamma| \leq T_a} (1 - H_\beta(v)) = N(v, T_a)$ . Hence,

$$\begin{aligned}
 & \left| \sum_{\rho \in \mathbb{Z}: |\gamma| \leq T_a} \frac{x^\rho - y^\rho}{\rho} \right| \\
 & \leq \frac{x-y}{y} \left[ y^{1/2} N(T_a) + \log y \int_{1/2}^{h(T)} y^v N(v, T_a) \, dv \right] \\
 (5.12) \quad & \triangleq \frac{2A(x-y)T \log T}{y} \left( y^{1/2} + T \log y \int_{1/2}^{h(T_a)} \left( \frac{y}{T^2} \right)^v \, dv \right) \\
 & = \frac{2A(x-y)T \log T}{y} \left( y^{1/2} + \frac{T \log y}{\log y - 2 \log T} \frac{y^v}{T^{2v}} \Big|_{v=1/2}^{h(T_a)} \right) \\
 & \leq \frac{2A(x-y)T^{2(1-h(T))} \log y \log T}{y^{1-h(T)} (\log y - 2 \log T)},
 \end{aligned}$$

recalling (5.10). In the last step, we have used the assumptions  $T \geq 14$  and  $T < y^{1/2}$  and the fact that  $\log y \geq \log y - 2 \log T$ . Also,  $T < T_a$  (more details).

Conclude from (5.8) with (5.9), (5.11), and (5.12) we obtain (3.8).

## 6. DIFFERENCES AS ERROR TERMS

*Right now, when we tackle problems without knowing the truth of the Riemann hypothesis, it's as if we have a screwdriver. But when we have it, it'll be more like a bulldozer. [24]*

Peter C. Sarnak

For the difference  $\varpi_2(x) - \varpi_2(y)$ , we write

$$\begin{aligned}
 & \varpi_2(x) - \varpi_2(y) = S_1 + S_2 + S_3 + S_4 + S_5 \\
 (6.1) \quad & = \left( \sum_{n \leq y} + \sum_{n=y} + \sum_{y < n < x} + \sum_{n=x} + \sum_{n > x} \right) \left( \delta_E\left(\frac{x}{n}\right) - \delta_E\left(\frac{y}{n}\right) \right) (\Lambda(n) - 1)
 \end{aligned}$$

where  $S_j$ 's for  $j \in \{1, 2, 3, 4, 5\}$  are defined in order, respectively.

We give estimates for  $S_2$  and  $S_4$  first. Similarly to (2.1), for  $v \in \mathbb{R}$  we define

$$(6.2) \quad \text{Id}_{\mathbb{N}}(v) = \begin{cases} 1, & \text{if } v \text{ is a natural number;} \\ 0, & \text{if } v \text{ is not a natural number.} \end{cases}$$

Recall the definition of  $\varpi_2(x)$  in (3.7) and that of  $\delta_E(v)$  in (5.3). One has

$$\begin{aligned}
 (6.3) \quad & S_2 = \left( \delta_E\left(\frac{x}{y}\right) - \delta_E(1) \right) \text{Id}_{\mathbb{N}}(y) (\Lambda(y) - 1) \\
 & = \left( \frac{1}{2\pi i} \int_{\mathcal{L}} \frac{x^s}{s y^s} \, ds - \frac{1}{\pi} \int_{T_a/m}^{\infty} \frac{du}{1+u^2} \right) \text{Id}_{\mathbb{N}}(y) (\Lambda(y) - 1) \triangleq O(T \log x).
 \end{aligned}$$

In the last step, we have used that  $|x^s/y^s| \leq 2$  for  $\frac{1}{2} < \sigma \leq 1$ ,  $|s| > \frac{1}{2}$  for  $s \in \mathcal{L}$ ,  $\Lambda(y) \leq \log y < \log x$ ,  $1 + u^2 > u^2$ , and  $T_a < T + 1$ . Similarly, one gets

(6.4)

$$\begin{aligned} S_4 &= \left( \delta_E(1) - \delta_E\left(\frac{y}{x}\right) \right) \text{Id}_{\mathbb{N}}(x)(\Lambda(x) - 1) \\ &= \left( \frac{1}{\pi} \int_{T_a/m}^{\infty} \frac{du}{1+u^2} - \frac{1}{2\pi i} \int_{\mathcal{R}} \frac{y^s}{sx^s} ds \right) \text{Id}_{\mathbb{N}}(x)(\Lambda(x) - 1) \triangleq O(T \log x), \end{aligned}$$

with  $y/x > 1/2$  and  $\sigma < 1$  for  $s \in \mathcal{R}$ .

To estimate  $S_5$ , we describe a technique dealing with integrals of complex functions along different routes. Let  $s_0 \in \mathbb{C}$  and  $s' \in \mathbb{C}$  be two different points on the complex plane and let  $\mathcal{C}_1$  and  $\mathcal{C}_2$  be two simple curves from  $s_0$  to  $s'$ . We assume that the union  $\mathcal{C}$  of  $\mathcal{C}_1$  and  $-\mathcal{C}_2$ , where  $-\mathcal{C}_2$  is the same route as  $\mathcal{C}_2$  along the opposite direction, is a simple closed curve. We note that  $s = 0$  is the unique pole of the integrand in the following integration. For any fixed  $x \in \mathbb{R}$ , we have

$$(6.5) \quad \int_{\mathcal{C}_1} \frac{x^s}{s} ds - \int_{\mathcal{C}_2} \frac{x^s}{s} ds = 2\pi i \delta$$

by the argument principle. In (6.5), the value of  $\delta = -1, 0$ , or  $1$ , equal to the index of  $s = 0$  with respect to  $\mathcal{C}$ . Suppose that  $F(s)$  is any uniquely defined regular function of  $s$  in an open region containing  $\mathcal{C}$ . Similarly to (6.5), we acquire

$$(6.6) \quad \int_{\mathcal{C}_1} F(s) \frac{x^s - y^s}{s} ds = \int_{\mathcal{C}_2} F(s) \frac{x^s - y^s}{s} ds,$$

as the constant term  $2\pi i \delta$  is canceled. Utilizing this equality, we transform  $S_5$  into an expression involving the integral along  $\mathcal{L}$  instead of  $\mathcal{R}$ . Recall that the union of  $-\mathcal{R}$  and  $\mathcal{L}$  is a simple closed curve whether the index of  $s = 0$  with respect to the union of  $-\mathcal{R}$  and  $\mathcal{L}$  is equal to  $0$ . By (6.6), one has

$$\begin{aligned} S_5 &= \sum_{n>x} \frac{\Lambda(n) - 1}{2\pi i} \int_{\mathcal{R}} \frac{\left(\frac{x}{n}\right)^s - \left(\frac{y}{n}\right)^s}{s} ds \\ &= \frac{1}{2\pi i} \sum_{n>x} (\Lambda(n) - 1) \int_{\mathcal{R}} \frac{1}{n^s} \cdot \frac{x^s - y^s}{s} ds \\ (6.7) \quad &= \frac{1}{2\pi i} \sum_{n>x} (\Lambda(n) - 1) \int_{\mathcal{L}} \frac{1}{n^s} \cdot \frac{x^s - y^s}{s} ds \\ &= \frac{1}{2\pi i} \int_{\mathcal{L}} \frac{x^s - y^s}{s} \sum_{n>x} \frac{\Lambda(n) - 1}{n^s} ds, \end{aligned}$$

recalling (5.3). This result is valid as long as the last series is convergent.

As for  $S_1$ , we interchange the order of the integration and summation. It results

$$\begin{aligned}
 S_1 &= \frac{1}{2\pi i} \sum_{n < y} \frac{\Lambda(n) - 1}{2\pi i} \left( \int_{\mathcal{L}} \frac{\left(\frac{x}{n} - \left(\frac{y}{n}\right)^s\right)}{s} \, ds \right) \\
 (6.8) \quad &= \frac{1}{2\pi i} \int_{\mathcal{L}} \frac{x^s - y^s}{s} \cdot \sum_{n < y} \frac{\Lambda(n) - 1}{n^s} \, ds.
 \end{aligned}$$

Also, the sum  $S_3$  may be rewritten as

$$\begin{aligned}
 S_3 &= \frac{1}{2\pi i} \sum_{y < n < x} (\Lambda(n) - 1) \int_{\mathcal{L}} \frac{\left(\frac{x}{n}\right)^s}{s} \, ds \\
 (6.9) \quad &= \frac{1}{2\pi i} \int_{\mathcal{R}} \frac{x^s}{s} \cdot \sum_{y < n < x} \frac{\Lambda(n) - 1}{n^s} \, ds.
 \end{aligned}$$

Combining (6.8), (6.9), and (6.7), we obtain that

$$(6.10) \quad S_1 + S_3 + S_5 = \frac{1}{2\pi i} \int_{\mathcal{L}} \frac{x^s - y^s}{s} \cdot \sum_{n \neq y \text{ and } n \neq x} \frac{\Lambda(n) - 1}{n^s} \, ds.$$

Concluding from (6.1) with (6.3), (6.4), and (6.10), we obtain

$$(6.11) \quad \varpi_2(x) - \varpi_2(y) = \frac{1}{2\pi i} \int_{\mathcal{L}} \mathcal{Z}(s) \frac{x^s - y^s}{s} \, ds + O(\log x).$$

Then, (3.9) follows.

## 7. ACKNOWLEDGEMENTS AND AFTERWORDS

*The greatest problem for mathematicians now is probably the Riemann hypothesis. But it's not a problem that can be simply stated.*

Andrew J. Wiles

It is my sincere wish to express my gratitude to Carl B. Pomerance for his prompting me into this direction. I would like to thank many of my colleagues for their encouragements and/or comments before, during, and/or after, the writing of this article and related, especially to Ronald L. Graham, D. Roger Heath-Brown, Glenn J. Fox, Andrew M. Odlyzko, Helmut Maier, Ju-Ping Wang, Andrew J. Granville, Hou-Rong Qin, Aloysius Helminck, Gong-Bao Li, Ze-Hua Chen, George E. Andrews, Chuan-Sheng Wu, Sidney W. Graham, Warren C. Wogen, Rashad M. Abrarov, Patrick B. Eberlein, and Xing-De Dai. Also, I want to thank the reviewers for their careful reading.

As direct consequences of Theorem 2, we now have many assertions being justified. For example, the Lindelöf hypothesis is valid. Another is about

the representation for the reciprocal of the Riemann zeta function. The series in the following is convergent for  $\sigma > \frac{1}{2}$ . Hence,

$$(7.1) \quad \frac{1}{\zeta(s)} = \sum_{n=1}^{\infty} \frac{\mu(n)}{n^s}, \quad \text{if } \sigma > \frac{1}{2},$$

where  $\mu(n)$  is the Möbius function. Let

$$(7.2) \quad \xi(s) = \pi^{-\frac{s}{2}} \xi_G(s) \xi_Z(s),$$

with

$$\xi_G(s) = \frac{s}{2} \Gamma\left(\frac{s}{2}\right) \quad \text{and} \quad \xi_Z(s) = (s-1) \zeta(s).$$

The next one is concerning with the  $\xi$  function defined in (7.2). We have

$$(7.3) \quad \Re \frac{\xi'(s)}{\xi(s)} > 0,$$

see [1]. In group theory, whenever  $n$  is sufficiently large, we have

$$(7.4) \quad \log g(n) < \frac{1}{\sqrt{\text{Li}(n)}},$$

where  $g(n)$  is the maximal order of elements in the symmetric group  $S_n$  of degree  $n$ , see [1] again. Also, the estimate on Titchmarsh's function

$$(7.5) \quad \arg \zeta(s) \Big|_{1+it}^{\lambda+it} = O\left(\frac{\log t}{\log \log t}\right), \quad \text{for } \frac{1}{2} \leq \lambda \leq 1,$$

is justified.

We mention two results in number theory here, which are not approximately but exact ones from the Riemann hypothesis. Robin showed in [32] that

$$(7.6) \quad \sigma(n) < e^\gamma n \log \log n,$$

for  $n \geq 5040$ . Lagarias proved in [27] that

$$(7.7) \quad \sigma(n) \leq H_n + e^{H_n} \log H_n,$$

for  $n \geq 1$  with the equality only for  $n = 1$ , where

$$(7.8) \quad H_n = 1 + \frac{1}{2} + \frac{1}{3} + \dots + \frac{1}{n}.$$

In passing, let me mention that any results under the Riemann hypothesis are valid. For instance, the existence of Mills' constant is validated now, see [5]. In the sense of Voronin's universality theorem,  $\zeta(s)$  approximates itself uniformly. For references, see [37], [31], or [23]. As a final point, we mention that one may revise the technique in this work to study the generalized Riemann hypothesis for the Dirichlet  $L$ -functions and the prime number theorem in arithmetic progressions with some substantial more work, see [7].

To conclude this article, let me say that some difficult-look problems in mathematics may not be so tough to deal with if we somehow find a neat way to tackle it. This is one of those examples, one may also see [7], [10], and, [11].



## REFERENCES

1. Borwein, P., Choi, S., Rooney, B., and Weirathmueller, A., *The Riemann hypothesis*, Springer, Berlin, Heidelberg, New York, Hong Kong, London, Milan, Paris, Tokyo, June 29, 2007.
2. Enrico Bombieri, *Prime territory: Exploring the infinite landscape at the base of the number system*, The Sciences, pp. 30–36, 1992.
3. Brent, R. P., *On the zeros of the Riemann zeta function in the critical strip*, Math. Comp. 33 (1979), no. 148, 136–1372.
4. Cipra, B., *A prime case of chaos*, What's Happening in the Mathematical Sciences, 4, pp. 2–17, 1999.
5. Cheng, Y. and Caldwell, C., *Determining Mills' Constant and a Note on Honaker's Problem*, Integer sequences, Article 05.4.1, Vol. 8, 2005.
6. Cheng, Y. and Fox, G., *Consequence from the remainder term of the prime number theorem*, to be submitted to Journal of Number Theory, In preparation, 2009.
7. Cheng, Y., Fox, G. J. and Graham, S. W., *Proof of the generalized Riemann hypothesis*, Initiated, 2009.
8. Cheng, Y. and Graham, S. W., *Estimates on the Riemann zeta function*, Rocky Mountain Journal of Mathematics, (4) 41, pp. 1261–1280, 2004.
9. Chandrasekharan, K., *Arithmetical Functions*, Springer-Verlag, Berlin, Heidelberg, New York, 1970.
10. Cheng, Y., *On primes beyond the Riemann hypothesis*, In preparation, 2009.
11. Cheng, Y., *Proof of the density hypothesis*, Submitted to the Journal of the American Mathematical Society, August 3, 2009.
12. Cheng, Y., *Estimates on primes between consecutive cubes*, Rocky Mountain Journal of Mathematics, to appear, 2009.
13. Cheng, Y., *Explicit estimates on prime numbers*, Rocky Mountain Journal of Mathematics, to appear, 2009.
14. Conrey, J.B., *More than two fifths of the zeros of the Riemann zeta function are on the critical line*, J. reine angew. Math. 399 (1989), 126.
15. Crandall R. and Pomerance C., *Prime Numbers – A computational perspective*, Springer, 2001.
16. Davenport, H., *Multiplicative Number Theory*, 2nd edition, Springer-Verlag, 1980.
17. Edward, H. M., *Riemann's zeta function*, Academic, New York, London, 1974.
18. Ford, K., *Zero-free regions for the Riemann zeta function*, Number Theory for the Millennium (Urbana, IL, 2000), II, pp. 25–56, 2002.
19. Gram, J. P., *Note sur les zeros de la fonction de Riemann*, Acta Math. 27 (1903), 289–304.
20. Gourdon, X., *The  $10^{13}$  first zeros of the Riemann zeta function, and zeros computation at very large height*, <http://numbers.computation.free.fr/Constants/Miscellaneous/zetazeros1e13-1e24.pdf>
21. Hoheisel, G., *Primzahlprobleme in der Analysis*, Sitz. Preuss. Akad. Wiss., 33, pp. 580–588, 1930.
22. Ivić, A., *The Riemann zeta function – Theory and Applications*, Dover Publications, Inc., Mineola, New York, 1985.
23. Karatsuba, A. A. and Voronin, Sergei M., *The Riemann Zeta-Function*, Walter de Gruyter, 1992.
24. Klarreich, E., *Prime time*, New Scientist, no. 2264, 2000.
25. Lu, Chang-Hai, *The Riemann hypothesis* [http://www.changhai.org/articles/science/mathematics/riemann\\_hypothesis/](http://www.changhai.org/articles/science/mathematics/riemann_hypothesis/).
26. Levinson, N., *More than one-third of the zeros of the Riemann zetafunction are on  $s = 1/2$* , Adv. Math. 13 (1974), 383–436.
27. Lagarias, J. C., *An elementary problem equivalent to the Riemann hypothesis*, Amer. Math. Monthly, 109, pp. 534–543, 2002.

28. Odlyzko, A. M. and Schnhage, A., *Fast algorithms for multiple evaluations of the Riemann zeta function*, Trans. Amer. Math. Soc. 309 (2): 797809
29. <http://planetmath.org/encyclopedia/ValueOfTheRiemannZetaFunctionAtS0.html>
30. Riemann, B., *Über die Anzahl der Primzahlen unter eine gegebener Grösse*, Monatsber. Akad., Berlin, pp. 671–680, 1859.
31. Bagchi, B., *Recurrence in Topological Dynamics and the Riemann hypothesis*, Acta Math. Hungar. 50, pp. 227–240, 1987.
32. Robin, G., *Grandes valeurs de la fonction somme des diviseurs et hypothèse de Riemann*, J. Math. Pures appl., **63**, pp. 187–213, 1984.
33. Selberg, A., *On the zeros of the zeta-function of Riemann*, Der Kong. Norske Vidensk. Selsk. Forhand. 15 (1942), 5962; also, Collected Papers, Springer-Verlag, Berlin - Heidelberg - New York 1989, Vol. I, 156–159.
34. Schoenfeld, L., *Sharper bounds for the Chebyshev functions  $\vartheta(x)$  and  $\psi(x)$  II*, Math. Comp., **30**, pp. 337–360, 1976. Corrigendum, *ibid.* **30**, pp. 900, 1976.
35. Turán, P., *On the remainder term of the prime number theorem formula II*, Acta Math. Acad. Sci. Hung. **1** (1950), pp 155–166 .
36. Turing, A. M., *A method for the calculation of the zeta-function*, Proc. London Math. Soc. (2) **48** (1943), 180–197.
37. Voronin, Sergei M., *Theorem on the Universality of the Riemann Zeta Function*, Izv. Akad. Nauk SSSR, Ser. Matem. 39, pp. 475–486, 1975. Reprinted in Math. USSR Izv. 9, pp. 443–445, 1975.

---

Drafted: October 04, 2007 – November 22, 2007.

Revised: March 23, 2009 – June 18, 2009.

Submitted: August 5, 2009.

Published: July ??, 2010.

---

*Keywords:* Prime numbers, the Riemann zeta function, the Riemann hypothesis, the prime number theorem, the density hypothesis, induction on the interval, differences, and the Lindelöf hypothesis.

---

ABSTRACT. The Riemann zeta function is a meromorphic function on the whole complex plane. It has infinitely many zeros and a unique pole at  $s = 1$ . Those zeros at  $s = -2, -4, -6, \dots$  are known as trivial zeros. The Riemann hypothesis is a conjecture of Riemann, from 1859, claiming that all nontrivial zeros of  $\zeta(s)$  lie on the line  $\Re(s) = \frac{1}{2}$ . Let  $x \geq 2$ . Define  $\Lambda(n) = \log p$  whenever  $n = p^m$  for a prime number  $p$  and a positive integer  $m$ , and zero otherwise. The Riemann hypothesis is then equivalent to the  $\varpi$ -form of the prime number theorem as  $\varpi(x) = O(x^{1/2} \log^2 x)$ , where  $\varpi(x) = \sum_{n \leq x} (\Lambda(n) - 1)$  with the sum running through the set of all natural integers. Let  $\mathcal{Z}(s) = -\frac{\zeta'(s)}{\zeta(s)} - \zeta(s)$ . Similar to that in the literature, we use the classical integral formula for the Heaveside function in the form of  $H(x) = \int_{m-i\infty}^{m+i\infty} \frac{x^s}{s} ds$  where  $m > 0$ , and  $H(x)$  is 0 when  $x < 1$ ,  $\frac{1}{2}$  when  $x = 1$ , and 1 when  $x > 1$ . However, we diverge from the literature by applying Cauchy's residue theorem to the function  $\mathcal{Z}(s) \cdot \frac{x^s}{s}$ , rather than  $-\frac{\zeta'(s)}{\zeta(s)} \cdot \frac{x^s}{s}$ , so that we may utilize the formula for  $m < 1$ , under certain conditions. Starting with the estimate on  $\varpi(x)$  from the known zero-free region of  $\mathcal{Z}(s)$ , we use induction to reduce the size of the exponent  $\theta$  in  $\varpi(x) = O(x^\theta \log^2 x)$ , while we also use induction on  $x$  when  $\theta$  is fixed. With the help of *the strong density hypothesis*, proved recently, we validate the aforementioned  $\varpi$ -form of the prime number theorem. Therefore, the Riemann hypothesis is proved.